

Conference Paper

Ecological and Geographical Gis-Analysis of Anomalies in Amphibians of Ukraine

V.M. Tytar¹, O.D. Nekrasova¹, and A.Yu. Marushchak²

¹I.I. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, st. Bogdana Khmel'nitskogo, 15, Kyiv, 01030, Ukraine

²2nd Year Master student at the department of zoology, ESC Institute of biology, Taras Shevchenko national university of Kyiv

Abstract

GIS modeling of 138 georeferenced point data of anomalies in amphibians in Ukraine (data from both the literature and personal field surveys) revealed a rather poor, but statistically significant correlation of the numbers of anomalies with the anthropogenic impact measured by the Human Footprint ($r=0,268$, $p=0,0025$). We consider that in certain cases the abundance of anomalies in amphibians could be used for purposes of bioindication.

Keywords: anomalies in amphibians, GIS-analysis.

Corresponding Author:

O.D. Nekrasova

oneks22@gmail.com

Received: 23 January 2018

Accepted: 20 April 2018

Published: 3 May 2018

Publishing services provided by
Knowledge E

© V.M. Tytar et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Selection and Peer-review under the responsibility of the Amphibian and Reptiles Anomalies and Pathology Conference Committee.

1. Introduction

For natural ecosystems in general and for amphibian populations in particular, anthropogenic impact is a very important factor. This factor is multidimensional, as it has direct and mediated aspects of influence. The decline of amphibian populations and the destruction of their natural environment are related to direct aspects of anthropogenic impact. While the emergence of specific morphological malformations is related to mediated factors of anthropogenic influence, such as pollution and other negative transformations of natural habitat.

Scientists have discovered, that morphological abnormalities originate more often in amphibians than in other terrestrial vertebrates [1, 3, 5]. Herewith an occurrence of amphibians with abnormalities in natural populations up to the level of 1.5-5% is considered normal [4, 9]. Such malformations could be genetically caused or acquired during the lifetime. The observation, that in isolated urban populations of moor frogs and marsh frogs, where the increased impact of mutagens cause abnormalities to emerge more often, than in other populations in more preferable conditions, is evidence of the sensitivity of the species to the anthropogenic "burden" [13-15, 23]. But this is not the only environment, where higher malformational rates occur. For example

OPEN ACCESS

in agroecosystems there were also big quantity of abnormalities of intestines (including gonads) observed among hybrids and edible frogs [13]. Pesticidal pollution of water reservoirs that leads to physiological and morphological changes in populations of green frogs, does also have a direct impact on their genome [24]. Hydrochemical water parameters (for example pH) have been found to influence the survival rate of amphibians and their morphogenesis in combination with other factors. In particular by forming of various morphological malformations [7].

2. Methods

To get a full picture of the effects of different possible factors on amphibian populations in the Ukrainian territory we used GIS-modeling and data visualization based on various indicators and factors found online [22]. In the SAGA GIS (version 2.2.5) package 138 georeferenced records of anomalies in amphibians (based on original and literature data - Marushchak, Muravynets, 2013; Fedoniuk, 2008; Kurtiak, 2010; Mik- itinets, 2014) have been used to extract the information contained in GIS environmental layers reflecting the bioclimate, land use, anthropogenic impact etc. The extracted information was kept in GIS layers. We got a 138X40 matrix, where the percentage of abnormalities in each sample of amphibians was considered. This information was analyzed in Statistica 8 pack using the method of multiple regression. The percentage of abnormalities was regarded as a dependent variable, and the other 39 - as independent.

We also used the bioclimatic parameters that are derived from the data on average monthly temperatures and precipitation. These variables are often used for modeling ecological niches, for example, when using such algorithms as the BIOCLIM (Busby, 1991; [22]).

We analyzed the confinement of habitats by comparing malformation levels in each area to the degree of influence of anthropogenic factors. As the initial parameters of anthropogenic burden the data from the interactive global map of human impact index (Human Footprint, HF), created in the Columbia combined forces of Wildlife Conservation Society (WCS) and the International Information Centre for Geosciences (CIESIN) in 2004 (Wildlife Conservation Society & Center for International Earth Science Information Network, Columbia University, 2005) [20] were used. This raster combines the layers representing the demographic parameters, land use indicators, infrastructure and access to territories.

From the European system ESDAC (European Soil Data Centre), which provides the European Union with all necessary soil information, we took GIS-maps showing the spreading of different soil types (max – 28 types) and levels of soils' pH.

For quantitative assessment of plant covering during the vegetation season (April–September) we used the normal derivative vegetation index (NDVI). This is an indicator for photosynthetic active biomass.

Additionally special metrics were applied to textural pictures of plant layers to understand the heterogenic patterns of the studied habitats [21].

3. Results

As a result of our work it can be concluded, that on huge territories it is hard to single out particular factors of one-way influence. Ecological problems and their impact on the occurrence of morphological abnormalities are best to be studied on a regional level. Alternatively classification principle could be used to combine territories and spheres of impact (ecological problems). Only for the index of anthropogenic influence (Human Footprint) we managed to show a particular impact on the variation of the dependent variable (the percentage of abnormalities in samples), although the correlation is fairly poor ($r=0.268$), however statistically significant ($p=0.0025$).

For a visual representation of the spread of amphibian abnormalities in Ukraine, using the spatial interpolation method (multilevel B-spline interpolation) in SAGA GIS, a map was compiled, where different colors correspond to the estimated percentage of abnormalities in amphibians (Figure 1).

4. Conclusion

It was concluded that, despite the fact that the nature of the abnormalities may be different and have complicated patterns, it can also be associated with a set of factors that are caused by human impact. It can be seen in relation to the index of human impact (Human Footprint) correlating weakly ($r = 0.268$), but significantly ($p = 0.0025$) with the percentage of malformations. This underlines the possible impact of different groups of anthropogenic factors, physical, chemical, biological, etc., acting on amphibian populations as multiple stressors, even an increase in dragonfly predators due to warmer temperatures [19]. Most likely amphibians are being subjected to a variety of human-induced impacts that are related to water quality. Special attention must be given to the presence of pollutants, pathogens, and global environmental changes that




Figure 1: The spatial interpolation model showing the frequency of abnormalities' occurrence in amphibians of Ukraine, compiled with the help of SAGA GIS program.

may affect amphibian growth and development, increase mortality, and eventually lead to population declines [2].

In our opinion, from a methodical standpoint the occurrence of abnormalities in amphibians may be used for bioindication and identification of adverse trends in the deterioration of the environment.

Since environmental factors have a strong spatial component, GIS is ideally suited to investigating stressors that might be responsible and hence it is a powerful analytical tool to understand the causes of amphibian population declines in particular environmental settings and stimulate further research [18].

Acknowledgement

To Ruben J. Teschner for help with the English translation, Oskyrko O. S, Muravynets O. A., Davidenko S. V., Stetsun G. A., Chernukha A. O. for helping to collect the samples.

References

- [1] Alford RA, Dixon MP & Pechmann KH): Global amphibian population declines. *Nature*, 2001; 412: 499–500.

- [2] Blaustein AR, David BW: The Puzzle of Declining Amphibian Populations. *Scientific American*, 1995; 272 (4): 52-57.
- [3] Blaustein AR, Han BA, Relyea RA, Johnson PT, Buck JC, Gervasi SS, & Kats LB: The complexity of amphibian population declines: understanding the role of cofactors in driving amphibian losses. *Annals of the New York Academy of Sciences* 1223.1; 2011: 108-119.
- [4] Borkin LJ, Bezman-Moseyko OS and Litvinchuk SN: Evaluation of animal deformity occurrence in natural populations (an example of amphibians). *Trudy Zoologicheskogo instituta RAN*, 2012; 316(4): 324-343 [in Russian].
- [5] Dubois A: Anomalies and mutations of natural populations of the *Rana "esculenta"* complex (Amphibia, Anura). *Mitteilungen aus dem Zoologischen Museum in Berlin*, 1979; 55: 59-87.
- [6] Fedoniuk OV: Amphibians and reptiles, their morphological anomalies and non-standard phenomena as consequence anthropological influences upon forestry ecosystems in Lviv region. In: *Interdepartmental Sci.-Tech. Misc.: placeForest, paper and wood industry*. 2008; 32: 355-360 [In Ukrainian].
- [7] Flyaks NL: The influence of pH on the survival and development of tailless amphibians of Sakhalin. *Systematics and ecology of amphibians and reptiles: Proceedings of ZIN AN SSSR*, 1986; 157: 152-166 [in Russian].
- [8] Flyaks NL and Borkin LJ: Morphological abnormalities and heavy metal concentrations in anurans of contaminated areas, eastern Ukraine. *Applied Herpetology*, 2004; 1: 229-264 [in Russian].
- [9] Kovalenko EE: Mass anomalies of the limbs of tailless amphibians. *Zhurnal obshchei biologii*, 2000; 61(4): 412-427 [in Russian].
- [10] Kurtyak FF: Anomalies of development of faintnesses in unisex hybrid populations *Pelophylax klepton esculenta* (Linnaeus, 1758) (Amphibia, Anura, Ranidae) in the Transcarpathians Lowland. In: *Scientific Bulletin of the Uzhgorod University*. 2010; Biology; 28: 132-134 [In Ukrainian].
- [11] Marushchak OY, Muravynets AA: Morphological abnormalities in natural populations of anurans (Amphibia, Anura) in Ukraine. In: *Proceeding of the Ukranian Herpetological society*. 4. Kyiv, 2013: 87-94 [In Ukrainian].
- [12] Mikitinez GI: Geography occurrence of in populations morphological abnormalities anura of steppe zone Ukraine. In: *Proceeding of international conference "Anomalies and Pathologies of Amphibians and Reptiles: Methodology, Evolutionary Impact, Possibilities for Estimation of Environmental Health"* (23-26 September 2013, Yekaterinburg, Russia): 144-149 [in Russian].

- [13] Nekrasova OD: Population structure and hybridization of green frogs of the "*Rana esculenta*" complex from urban territories of Middle Dnieper basin. - Manuscript. Thesis for submitting to the degree of Phd. Speciality: 03.00.02. - Zoology. - Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine, Kyiv, 2002: 21 [in Russian].
- [14] Nekrasova OD, Mezhzherin SV, Morozov-Leonov SYu, Sytnik YuM: A cases of mass polymelia in the lake frog (*Rana ridibunda* Pall., 1771) from Kyiv. Scientific Bulletin of the Uzhgorod University, (Ser. Biol.), 2007: 92-95 [in Russian].
- [15] Nekrasova OD: Classification of Amphibians Anomalies. In: Proceeding of the Ukrainian Herpetological society. 1; Kyiv, 2008: 55-58 [in Russian].
- [16] Nekrasova OD: Some aspects of anomaly's manifestation in amphibians coloration. In: Proceeding of international conference "Anomalies and Pathologies of Amphibians and Reptiles: Methodology, Evolutionary Impact, Possibilities for Estimation of Environmental Health" (23-26 September 2013, Yekaterinburg, Russia): 144-149 [in Russian].
- [17] Ouellet M, Bonin J, Rodrigue J, DesGranges JL, Lair S: Hindlimb deformities (ectromelia, ectrodactyly) in free-living anurans from agricultural habitats. J. Wildl Dis; 1997; 33: 95-104.
- [18] Rao M, Focht W, Bantle JA: Application of GIS in Environmental Policy Analysis. Oklahoma Politics, 2001; 10: 141-155.
- [19] Reeves M, Jensen P, Dolph C, Holyoak M, Trust K: Multiple stressors and the cause of amphibian abnormalities. Ecological Monographs, 2010; 80(3): 423-440.
- [20] Sanderson EW, Jaiteh M, Levy MA, Redford KH, Wannebo AV, & Woolmer G: The Human Footprint and the Last of the Wild: The human footprint is a global map of human influence on the land surface, which suggests that human beings are stewards of nature, whether we like it or not. BioScience, 2002; 52(10): 891-904.
- [21] Tuanmu M-N and Jetz W: A global, remote sensing-based characterization of terrestrial habitat heterogeneity for biodiversity and ecosystem modeling. Global Ecology and Biogeography. 24(11); November 2015: 1329-1339.
- [22] Tytar VM: Analysis of home ranges in species: an approach based on modeling the ecological niche. Vestnik Zoologii. suppl. 2011; 25: 93 [In Ukrainian].
- [23] Vershinin VL: Morphological abnormalities in urban amphibians. Ecology. Sverdlovsk, 1989; (3): 58-66 [in Russian].
- [24] Zhukova TI, Kubantsev BS, Burlachenko TL: Some reactions of populations of lake frog on pesticide pollution of water reservoirs. In: Proceedings of scientific.

Anthropogenic impact on the populations of the animals. Volgograd, 1986: 61-81 [in Russian].